



HAL
open science

Climate Change and social transformations in the past (12ka BP): from field data acquisition towards socio-ecological modeling

Laurent Lespez, Maria-Angela Bassetti, Jean-François Berger, Jean-Michel Carozza, Laurent Carozza, Nathalie Combourieu Nebout, Laurent Dezileau, Arthur Glais, Matthieu Ghilardi, Catherine Kuzucuoğlu, et al.

► To cite this version:

Laurent Lespez, Maria-Angela Bassetti, Jean-François Berger, Jean-Michel Carozza, Laurent Carozza, et al.. Climate Change and social transformations in the past (12ka BP): from field data acquisition towards socio-ecological modeling. Conférence MISTRALS PALEOMEX, Oct 2017, Montpellier, France. 2016. hal-01683548

HAL Id: hal-01683548

<https://univ-tlse2.hal.science/hal-01683548v1>

Submitted on 18 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.



L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Laurent Lespez¹, Maria-Angela Bassetti², Jean-François Berger³, Jean-Michel Carozza⁴, Laurent Carozza⁵, Nathalie Combourieu-Nebout, Laurent Dezileau⁶, Arthur Glais⁷, Matthieu Ghilardi⁸, Catherine Kuzucuoglu¹, Odile Peyron⁹, Pierre Sabatier¹⁰, Mehdi Saqalli⁴, Boris Vannière¹¹, Marie-Alexandrine Sicre¹², Bassem Jalali¹³, and Paleomex team

1. LGP-UMR CNRS 8591 - Université de Paris Est - Créteil et Université de Paris 1, 2. CEFREM-UMR CNRS 5110, 3. EVS-UMR CNRS 5600 - Université Lyon 2, 4. LIENS, UMR 7266, CNRS - Université de La Rochelle, 5. GEODE-UMR CNRS 5602 - Université Toulouse 2 « Jean Jaurès », 6. Géosciences Montpellier UMR CNRS 5243, 7. LETG Caen UMR CNRS 6554 - Université de Caen Normandie, 8. CEREGE-UMR CNRS 7330 - Université d'Aix-Marseille, 9. ISEM Montpellier, 10. Laboratoire EDYTEM - UMR CNRS 5204, Université Savoie Mont Blanc, 11. Chrono-Environnement-UMR CNRS 6249, Université de Franche-Comté, Besançon, 12. LOCEAN, Sorbonne universités, 13. GEOGLOB, Université de Sfax, Sfax, Tunisie.

Objectives and challenges

- “ Climatic trends in Mediterranean areas during the Holocene (from 12 ka BP)
- “ Definition of the spatial and temporal variability of the Rapid Climate Changes (RCCs)
- ⇒ Climate change and impact on cultural and political dynamic?
- Neolithic (9.2, 8.2 and 6-5 ka BP)
- Bronze Age (4.2 ka cal BP)
- Final Bronze Age and Historical periods (3.2-2.8 and 1.3 et 0.7 ka cal. BP)



Methods : 4 transects – multiproxies analyses




- Long marine sequences
- Analyses of long pollen and fire signature series for high resolution climate changes analyses (e.g. modern analogs - Peyron et al., 2017; Vannière et al., 2015)
- High resolution analyses of lake and fluvial sequences (ex. 8.2 tripartition in Berger et al., CP 2016)
- Socio-political changes : cultural areas, settlement, political changes (e.g. Carozza et al., 2015; Lespez et al., 2016a, b)

⇒ Climate modelisation
 ⇒ Climate-Environment-Societies interactions (ex. Saqalli, M., Tilman B. 2015.)

Paleomex in the Lion's Gulf

Improve climate and environmental change: seesaw across the Mediterranean basin




Sea surface temperatures and land-derived input time series were generated from the Gulf of Lions inner-shelf sediments (NW Mediterranean Sea) using alkenones and high-molecular-weight odd-carbon numbered n-alkanes. The SST record depicts three main phases: a warm Early Holocene (~18 ± 0.4 °C) followed by a cooling of ~3 °C between 7000 and 1000 BP, and rapid warming from ~1850 AD onwards.

Several superimposed multi-decadal to centennial-scale cold events of ~1 °C amplitude were also identified.

Concentrations show a broad increase from the Early Holocene towards the present with a pronounced minimum around 2500 BP and large fluctuations during the Late Holocene.

Sediments of the inner shelf seems to be originated mainly from the Upper Rhone River watershed, primarily delivered during positive North Atlantic Oscillation (NAO).

Conceptual model of Climate/Environment/Society interactions




4.2 ka BP climatic event and settlement pattern changes from the Late Neolithic to the Early Bronze Age in western Mediterranean?

- Effects of RCC lasting 3-4 centuries around the 4.2 ka BP event, c. 2.2 ka BC recorded in the lake, fluvial and soil systems
- A temporal tripartite structure with 2 wet periods in Southern France
- Change in the human settlement system around 2.3-2.2 ka BC:
 - In lowland areas, the number of settlements decreased significantly along the river systems during a period of very high hydrosedimentary discharges, dryness, and fire activity.
 - Environmental changes (glacial retreat) permitted the exploitation of copper ore resources at very high altitudes of above 2,400 m allowed for an exploitation of alpine copper, as in Saint-Véran (SE France) and archaeological findings have revealed a growth in human pressure in mountain areas, specifically in the Pyrenees (SW France).
- ⇒ Change of settlement from lowland area to mountainous areas may have resulted in a spatial reorganization at a regional level, but not in a global societal collapse.

Modelling Climate/Environment/Society interactions

Dynamic and spatially-explicit modeling is the only way for combining together biophysical elements such as climate, soils, and land cover, with archaeological and socio-anthropology/socio-economy based hypotheses in the functioning of the Neolithic societies.

However, Paleo-environmentalists provide climate and landscape reconstructions with a century-scale temporal precision while, to understand the consequences on rural populations, one should translate these data into the seasonal cropping and hunting/gathering activities; Archaeologists provide site-specific habitat and activity descriptions for specific time periods while, to extend such reconstructions for applying it on a regional level, the one occupied by the same culture, a generic and adaptable behavior rationality should be hypothesized, combining social constraining rules and production practices.



We propose a spatially explicit (scale 1ha) and temporally defined (scale: 1 season) multi-agent modelisation.

The GAMA platform gama-platform.org/ built in Toulouse (France) is the software reference for such a collective and long-term modelling process

References:
 Berger J. F., Lespez L., Kuzucuoglu C., Glais A., Hourani F., Barra A., Guillaune J., 2016. Early to Mid-Holocene Neolithic/RCC interactions to East Mediterranean basins (Greece, Turkey, Cyprus). *Climate of the Past*, 12, 1847-1877, 2016, <http://www.clim-past.net/12/1847/2016/>
 Carozza L., Berger, J. F., Burens-Carozza, A., Marcigny, C., 2015. Society and environment in Southern France from the 3rd millennium BC to the beginning of the 2nd millennium BC: 2200 BC a tipping point? 2200 BC a climatic breakdown as a cause for the collapse of the old world? In Tagungen des Landesmuseum für vorgeschichte, Halle, Band 12, p. 333-362.
 Ghilardi M., Lespez L., (eds.) 2017. Geoarchaeology of the Mediterranean Islands: from "lost worlds" to vibrant places, Special Issue *Journal of Archaeological Science Reports*.
 Lespez L., Glais A., López-Sáez J.-A., Le Dren V., Tárison Z., Davidson R., Brée L., Malamoudou D., 2016. Mid-Holocene rapid environmental changes and human adaptation in Northern Greece. *Quaternary Research* 85, 2, 227-244.
 Lespez L., Carozza L., Berger J. F., Kuzucuoglu C., Ghilardi M., Carozza J.-M., Vannière B., and the ArchoMed team, 2016. Rapid climatic change and social transformations: Uncertainties, Adaptability and Resilience. In Altieri (ed.) *The Mediterranean region under climate change. A scientific Update*, 10 p.
 Peyron O., Combourieu-Nebout, N., Brayshaw D., Goring, S., Andrieu-Ponel, V., Desprat, S., ... & Kotthoff, U. (2017). Precipitation changes in the Mediterranean basin during the Holocene from terrestrial and marine pollen records: a model-data comparison. *Climate of the Past*, 13(3), 249.
 Vannière B., Blarquez O., Rius D., Doyen E., Brucher T., Colombaroli D., Connor S., Feurdean A., Hickler T., Lemmen C., Leys B., Massa C., Olofsson J. (2016) 7000-year human legacy of elevation-dependent European fire regimes. *Quaternary Science Reviews* 132, 206-212.
 Jalali B., Sicre M.-A., Bassetti M.-A., Kallel N. 2016. Holocene climate variability in the North-Western Mediterranean Sea (Gulf of Lions). *Climate of the Past Discussions*, European Geosciences Union (EGU), 2016, pp.91-120.
 Sicre M.-A., Jalali B., Marttat B., Schmidt S., Bassetti M.-A., Kallel N. 2016. Sea surface temperature variability in the North-Western Mediterranean Sea (Gulf of Lions) during the Common Era. *Earth and Planetary Science Letters* 456, 124-133
 Saqalli, Baum. 2015. Pathways for scale reconciliation: Building ecological sociomodeling methodologies for a reconstruction of human past dynamics over a landscape. In: Chap. 9. Barceló J. A., Del Castillo F. (eds.) *Simulating prehistoric and ancient worlds. Springer Computational Social Sciences*. 233-255.